

THE CANADIAN MEDICAL ASSOCIATION

LE JOURNAL DE

L'ASSOCIATION MÉDICALE CANADIENNE

ORIGINAL ARTICLES

Polycystic Ovaries Associated with Congenital Adrenal Hyperplasia. <i>O. J. Lucis, R. Hobkirk, C. H. Hollenberg, S. A. MacDonald and P. Blahey</i>	1
A Follow-up Evaluation of the Treatment of Chronic Alcoholics. <i>G. Sereny and M. Fryatt</i>	8
Management and Surgical Treatment of Ruptured Intracranial Aneurysms. <i>P. D. Moyes</i>	13
Acute Purulent Meningitis in Alaska Natives: Epidemiology, Diagnosis and Prognosis. <i>R. Fortune</i>	19

CURRENT PROGRESS

Interferon: A Changing Picture. <i>R. P. B. Larke</i>	23
--	----

MEN AND BOOKS

Medicine and the Biblical World. <i>J. B. Hardie</i>	32
---	----

GENERAL PRACTICE

Cervical Traction in the Home. <i>C. A. Laurin</i>	36
---	----

CASE REPORT

Asthma Associated with the Use of Nitrofurantoin. <i>C. H. A. Walton</i>	40
---	----

EDITORIALS AND ANNOTATIONS

Rite of Passage	42
Taking Their Medicine	43

LETTERS TO THE JOURNAL

Semantics and "The Pill". <i>Anne C. Lowe</i>	44
<i>M. Nadasdi</i>	44
<i>R. A. Jacques</i>	44

THE LONDON LETTER

The Majority Say Yes; Adverse Reactions to Drugs; Psychiatric Disorders in the Aged; Effect of Car Radios on Driving. <i>S. S. B. Gilder</i>	46
---	----

ASSOCIATION NEWS

The Editor	47
------------------	----

NEWS AND VIEWS ON THE ECONOMICS OF MEDICINE

Collective Bargaining	51
-----------------------------	----

Obituaries	48	Instructions to Contributors	29
Book Reviews	50	Medical News in Brief	33
Abstracts	12	Index of Advertisers	36
Provincial News	24		



COMPOSITION: Meclizine hydrochloride 12.5 mg. and niacin 50 mg. per tablet.

USES: Antivert is indicated in the treatment of vertigo, whether due to cerebral arteriosclerosis, Ménière's syndrome, labyrinthitis, radiation sickness, vestibular dysfunction or other related conditions. Antivert has also been found valuable in the treatment of vascular headache.

DOSAGE: One Antivert tablet 3 times a day before meals. If excessive flushing occurs, tablets should be taken immediately after meals.

Patients should be made aware of the frequent, short-lived reactions experienced with niacin therapy: cutaneous flushing, sensations of warmth, tingling and itching, burning of the skin, increased gastrointestinal motility, and sebaceous gland activity. These reactions are coincidental to vasodilation produced by niacin. In explaining these reactions to the patient, it is suggested that they be regarded as desirable, being a physiological sign that the niacin is carrying out its intended function.

PRECAUTIONS: Meclizine may occasionally induce drowsiness. The physician should inform patients of the need for due precautions when engaged in activities requiring alertness.

Because of the vasodilation produced by niacin, severe hypotension and hemorrhage are contraindications to Antivert therapy.

SUPPLY: Scored tablets (blue), in bottles of 100.

ANTIVERT
stops
vertigo

JANUARY 1, 1966

Volume 94 • Number 1

Published every week by THE CANADIAN MEDICAL ASSOCIATION
L'ASSOCIATION MÉDICALE CANADIENNE

[COPYRIGHTED] 150 St. George Street, Toronto 5.

Subscription rates on advertising page 36

WOODS HOLE, MASS., U.S.A.
MARINE BIOLOGICAL LAB.